

DIAGNOSIS CODES QUICK REFERENCE GUIDE

This guide is designed to support the reimbursement process for both providers and payers by providing coding information for Entyvio (vedolizumab). Providers are responsible for determining and submitting the appropriate codes, charges, and modifiers for all medically appropriate services and products. Please contact individual payers for current and specific coding, coverage, and payment policies.

The following ICD-9-CM or ICD-10-CM diagnosis codes may be appropriate to describe these disease states:

ULCERATIVE COLITIS (UC) ICD-9 TO ICD-10 CONVERSION TABLE¹

ICD-9 diagnosis codes			ICD-10 diagnosis codes	
Code	Description		Code	Description
556.0	Ulcerative (chronic) enterocolitis	→	K51.80	Other ulcerative colitis without complications
556.1	Ulcerative (chronic) ileocolitis	→	K51.80	Other ulcerative colitis without complications
556.2	Ulcerative (chronic) proctitis	→	K51.20	Ulcerative (chronic) proctitis without complications
556.3	Ulcerative (chronic) proctosigmoiditis	→	K51.30	Ulcerative (chronic) rectosigmoiditis without complications
556.5	Left-sided ulcerative (chronic) colitis	→	K51.50	Left-sided colitis without complications
556.6	Universal ulcerative (chronic) colitis	→	K51.00	Ulcerative (chronic) pancolitis without complications
556.8	Other ulcerative colitis	→	K51.80	Other ulcerative colitis without complications
556.9	Ulcerative colitis, unspecified	→	K51.90	Ulcerative colitis, unspecified, without complications

CROHN'S DISEASE (CD) ICD-9 TO ICD-10 CONVERSION TABLE¹

ICD-9 diagnosis codes			ICD-10 diagnosis codes	
Code	Description		Code	Description
555.0	Regional enteritis of small intestine	→	K50.00	Crohn's disease of small intestine without complications
555.1	Regional enteritis of large intestine	→	K50.10	Crohn's disease of large intestine without complications
555.2	Regional enteritis of small intestine with large intestine	→	K50.80	Crohn's disease of both small and large intestine without complications
555.9	Regional enteritis of unspecified site	→	K50.90	Crohn's disease, unspecified, without complications

Please see Indications and Important Safety Information on next page.

Reference: 1. AAPC. ICD-10 Code Translator. <https://www.aapc.com/icd-10/codes>. Accessed September 8, 2015.

INDICATIONS: ENTYVIO (vedolizumab)

Adult Ulcerative Colitis (UC)

Adult patients with moderately to severely active UC who have had an inadequate response with, lost response to, or were intolerant to a tumor necrosis factor (TNF) blocker or immunomodulator; or had an inadequate response with, were intolerant to, or demonstrated dependence on corticosteroids:

- inducing and maintaining clinical response
- inducing and maintaining clinical remission
- improving endoscopic appearance of the mucosa
- achieving corticosteroid-free remission

Adult Crohn's Disease (CD)

Adult patients with moderately to severely active CD who have had an inadequate response with, lost response to, or were intolerant to a TNF blocker or immunomodulator; or had an inadequate response with, were intolerant to, or demonstrated dependence on corticosteroids:

- achieving clinical response
- achieving clinical remission
- achieving corticosteroid-free remission

IMPORTANT SAFETY INFORMATION

- ENTYVIO (vedolizumab) for injection is contraindicated in patients who have had a known serious or severe hypersensitivity reaction to ENTYVIO or any of its excipients.
- Infusion-related reactions and hypersensitivity reactions including anaphylaxis have occurred. Allergic reactions including dyspnea, bronchospasm, urticaria, flushing, rash, and increased blood pressure and heart rate have also been observed. If anaphylaxis or other serious allergic reactions occur, discontinue administration of ENTYVIO immediately and initiate appropriate treatment.
- Patients treated with ENTYVIO are at increased risk for developing infections. Serious infections have been reported in patients treated with ENTYVIO, including anal abscess, sepsis (some fatal), tuberculosis, salmonella sepsis, *Listeria* meningitis, giardiasis, and cytomegaloviral colitis. ENTYVIO is not recommended in patients with active, severe infections until the infections are controlled. Consider withholding ENTYVIO in patients who develop a severe infection while on treatment with ENTYVIO. Exercise caution in patients with a history of recurring severe infections. Consider screening for tuberculosis (TB) according to the local practice.
- Although no cases of PML have been observed in ENTYVIO clinical trials, JC virus infection resulting in progressive multifocal leukoencephalopathy (PML) and death has occurred in patients treated with another integrin receptor antagonist. A risk of PML cannot be ruled out. Monitor patients for any new or worsening neurological signs or symptoms. Typical signs and symptoms associated with PML are diverse, progress over days to weeks, and include progressive weakness on one side of the body or clumsiness of limbs, disturbance of vision, and changes in thinking, memory, and orientation leading to confusion and personality changes. If PML is suspected, withhold dosing with ENTYVIO and refer to a neurologist; if confirmed, discontinue ENTYVIO dosing permanently.
- There have been reports of elevations of transaminase and/or bilirubin in patients receiving ENTYVIO. ENTYVIO should be discontinued in patients with jaundice or other evidence of significant liver injury.
- Prior to initiating treatment with ENTYVIO, all patients should be brought up to date with all immunizations according to current immunization guidelines. Patients receiving ENTYVIO may receive non-live vaccines and may receive live vaccines if the benefits outweigh the risks.
- Most common adverse reactions (incidence $\geq 3\%$ and $\geq 1\%$ higher than placebo): nasopharyngitis, headache, arthralgia, nausea, pyrexia, upper respiratory tract infection, fatigue, cough, bronchitis, influenza, back pain, rash, pruritus, sinusitis, oropharyngeal pain, and pain in extremities.

Please click [here](#) to read the full [Prescribing Information](#), including [Medication Guide](#).

ENTYVIO is a trademark of Millennium Pharmaceuticals, Inc., registered with the U.S. Patent and Trademark Office, and is used under license by Takeda Pharmaceuticals America, Inc.

